

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶:

C07D 495/04, A61K 31/505, 31/44 // (C07D 495/04, 333:00, 239:00) (C07D 495/04, 333:00, 221:00)

(11) International Publication Number:

WO 99/24440

(43) International Publication Date:

20 May 1999 (20.05.99)

(21) International Application Number:

PCT/IB98/01691

A1

(22) International Filing Date:

22 October 1998 (22.10.98)

(30) Priority Data:

60/065,097

11 November 1997 (11.11.97) US

(71) Applicant (for all designated States except US): PFIZER PRODUCTS INC. [US/US]; Eastern Point Road, Groton, CT 06340 (US).

(72) Inventors; and

- (75) Inventors/Applicants (for US only): MUNCHHOF, Michael, John [US/US]; 266 West Road, Salem, CT 06420 (US). SOBOLOV-JAYNES, Susan, Beth [US/US]; 4 Piney Branch Road, Ivoryton, CT 06442 (US).
- (74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., 235 East 42nd Street, New York, NY 10017 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published .

With international search report.

(54) Title: THIENOPYRIMIDINE AND THIENOPYRIDINE DERIVATIVES USEFUL AS ANTICANCER AGENTS

(1)

(2)

(57) Abstract

The invention relates to compounds of formulas (1) and (2) and to pharmaceutically acceptable salts and hydrates thereof, wherein X¹, R¹, R² and R¹¹ are as defined herein. The invention also relates to pharmaceutical compositions containing the compounds of formulas (1) and (2) and to methods of treating hyperproliferative disorders in a mammal by administering the compounds of formulas (1) and (2).